

MORE FUN WITH FUNDAMENTAL IDENTITIES

Tips

- 1) If in doubt, change everything to sines & cosines.
- 2) Use identities which make terms cancel.
- 3) If fractions are added or subtracted, make common denominators.
- 4) Change both sides to the same trig functions, so you can see what you are trying to equal.
- 5) If you need an expression to contain squared terms, try multiplying by the conjugate.
- 6) If terms have powers > 2 , try to factor.

$$\frac{\cos^2 x + 3 \sin x - 1}{3 + 2 \sin x - \sin^2 x} = \frac{1}{1 + \csc x}$$

$$\frac{\cancel{x} - \sin^2 x + 3 \sin x - \cancel{1}}{3 + 2 \sin x - \sin^2 x} = \frac{1}{\frac{\sin x}{\sin x} + \frac{1}{\sin x}}$$

$$\frac{3 \sin x - \sin^2 x}{3 + 2 \sin x - \sin^2 x} = \frac{1}{\frac{\sin x + 1}{\sin x}}$$

$$\frac{\sin x (\cancel{3 - \sin x})}{(\cancel{3 - \sin x})(1 + \sin x)} = \frac{\sin x}{\sin x + 1}$$

$$\frac{1}{\sec x - \tan x} \cdot \frac{(\sec x + \tan x)}{(\sec x + \tan x)} \sec x + \tan x$$

$$\frac{\sec x + \tan x}{\sec^2 x - \tan^2 x} = \sec x + \tan x$$

Calculator

$$\frac{\cot x}{\cos x} - \frac{\csc^2 x}{\sec x} = \frac{\sin x - \cos x}{\sin^2 x}$$