Name		

ALGEBRA II JOURNAL Square Roots & Complex Numbers

1.	adding or subtracting square roots, they must contain values. when			
2.	(a) If the denominator of a fraction contains a single square root, it can be moved to the numerator by			
	(b) If the denominator of a fraction contains a square root plus another number $(a + \sqrt{b})$, it			
	can be moved to the numerator by multiplying by which is called			
	the (c) When a square root is moved from the denominator of a fraction to the numerator, the			
	(c) When a square root is moved from the denominator of a fraction to the numerator, the process is called			
3.	When solving an equation like $x^2 = 81$, you must			
	and remember to add to your answer.			
4.	Numbers that result from the square roots of negative numbers are called			
	numbers.			
5.	Complex numbers earn this name because they have two parts: and			
6.	ALL numbers are numbers.			
7.	The value of <i>i</i> to a very large power can be found by			
8.	Before performing any arithmetic operation (+, -, x, /) between the square roots of two negative numbers, you must first			
9.	If a fraction has only 4 <i>i</i> in the denominator, you would rationalize the denominator by			
10	. If a fraction has 2 + 5 <i>i</i> in the denominator, you would move it to the numerator by			
11	. (a) The Mandelbrot Set is an example of a whose primary characteristic is			
	(b) The Mandelbrot Set is created on a coordinate axis in which the <i>x</i> -axis is the			
	axis and the <i>y</i> -axis is the axis. (c) The Mandelbrot Set is created with the function $f(x) =$ through the			
	process of			
	(d) Extra Credit: Show 3 iterations of the function $f(x) = x^2 - 10$ beginning at $x = 1$ the last digit of your calculator number. (Example: If your calculator number is NC95, iterate using $x = 1$)			

12. Important Rules, Formulas, Etc. a) i =_____

a)
$$i =$$

b) List the 4 powers of *i* and the saying for how to remember them.